



## Size distribution measurement of fine and ultrafine particle emission from cooking activities

Evelyne Gehin, Olivier Ramalho, Séverine Kirchner

### ► To cite this version:

Evelyne Gehin, Olivier Ramalho, Séverine Kirchner. Size distribution measurement of fine and ultrafine particle emission from cooking activities. European Aerosol Conference 2008, Aug 2008, Thessaloniki, Greece. pp.T10A002O. hal-00701718

**HAL Id: hal-00701718**

**<https://hal.science/hal-00701718>**

Submitted on 25 May 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Size distribution measurement of fine and ultrafine particle emission from cooking activities

Evelyne Géhin<sup>a</sup>, Olivier Ramalho<sup>b</sup>, Severine Kirchner<sup>b</sup>

<sup>a</sup>Centre d'Etude et de Recherche en Thermique Environnement et Systèmes, Université Paris 12, Avenue du Général de Gaulle, 94010 Créteil, France

<sup>b</sup>Laboratoire de Microbiologie des Environnements Intérieurs, Centre Scientifique et Technique du Bâtiment, Champs sur Marne, France

Keywords: Emission factor, indoor particles, particle size distribution, measurement

The most commonly cited sources for particle emission indoor are: cigarette smoke, cooking, heating with fuel, fireplace, candles or incense burning, use of vacuum cleaner, use of spray (He et al., 2004 ; Hussein et al., 2006, Afshari et al., 2005). Some other sources are also quoted, like the laser printers, the heater and sauna heating. There is very few results on size number distribution emission rate in the range between 5 nm and 1 µm (fine and ultrafine particles) despite the fact that exposure to fine and ultrafine airborne particles has been identified as a factor that could affect human health (See and Balasubramaniana, 2006). We present here a comparison of particles emission rate (diameter in the range 5 nm to 1 µm) on 8 different cooking activities (cooking with electric stove or oven). All the necessary staff for the activities was purchased from common chain stores and the test procedure is most of possible representative of the usual way.

The experiments were carried out in the laboratory of the Centre Scientifique et Technique du Bâtiment (CSTB) in an hexagonal experimental chamber. An ascending current of filtered air is created in the chamber (average vertical air velocity of 10 cm.s<sup>-1</sup>). In order to create realistic conditions of sources emission all the cooking activities are done from outside of the chamber with airtight gloves. The particle size spectrometer used in this study is a DMS500 (Cambustion) which measure particle concentration as a function of the equivalent electrical mobility diameter (16 size fraction per decade) from 5 nm to 1000 nm at a maximum rate of 10 samples/second. The particle concentrations are measured every second for minimizing RMS noise.

Assuming that the air in the chamber is well-mixed the particle size balance equation of the particle number concentration at time  $t$  is described by:

$$\frac{\partial N_{d,t}}{\partial t} = -\frac{Q}{V} \times N_{d,t} - k_d^{dep} \times N_{d,t} + \frac{q_{d,t}^{emi} + J_{d,t}^{coag} + J_{d,t}^{cond}}{V}$$

Where  $Q$  is the ventilation rate in the chamber,  $N_{d,t}$  is the number concentration (cm<sup>-3</sup>) of particle with diameter between  $d$  and  $(d+dd)$ ,  $q_{d,t}^{emi}$  is a global emission rate (s<sup>-1</sup>) for particle with diameter between  $d$  and  $(d+dd)$  including emission rate of the source, nucleation, re-suspension, sinks.  $k_d^{dep}$  is the deposition rate of the particles on the surfaces of the experimental chamber.  $J_{d,t}^{coag}$  and  $J_{d,t}^{cond}$  are the

change rate of particles with diameter between  $d$  and  $(d+dd)$  due to coagulation and condensation effects. The emission rate is measured during all the cooking activity and a mean equivalent emission rate (MEER) is calculated by:

$$\overline{q_d^{equ}} = \frac{1}{(t_2 - t_1)} \times \int_{t_1}^{t_2} (q_{d,t}^{emi} + J_{d,t}^{coag} + J_{d,t}^{cond}) dt$$

The mean total decay rate ( $k_d = \frac{Q}{V} + k_d^{dep}$ ) is

determined experimentally and range between 0.0153 s<sup>-1</sup> and 0.0184 s<sup>-1</sup> with a minimum value for particles with a diameter of 350 nm.

The MEER of cooking activities present a mode around 20-40 nm with a maximum between 10<sup>9</sup> s<sup>-1</sup> and 10<sup>12</sup> s<sup>-1</sup>. Frying meat or fish lead to a total mean equivalent emission rate of (9±6)×10<sup>10</sup> s<sup>-1</sup> while cooking meat or fish in an oven lead to total mean equivalent emission rate of (9±4)×10<sup>10</sup> s<sup>-1</sup>. Cooking pasta or heat the stove will lead to total mean equivalent emission rate equal to (7±5)×10<sup>9</sup> s<sup>-1</sup>. These emission rates are comparable to what was observed by Afshari et al. (2005). They reported emission rates for particles with a diameter between 20 nm and 1 µm of 1.38×10<sup>10</sup> s<sup>-1</sup> for frying meat on an electric stove and 1.13×10<sup>10</sup> s<sup>-1</sup> when heating the electric stove.

This work was supported by the french « Ministère de l'écologie et du développement durable ».

Afshari, A., Matson, U., Ekberg, L.E., 2005.

Characterization of indoor sources of fine and ultrafine particles: a study conducted in a full-scale chamber, *Indoor Air*, 15, 141-150.

He, C., Morawska, L., Hitchins, J., Gilbert, D., 2004.

Contribution from indoor sources to particle number and mass concentrations in residential houses, *Atmospheric Environment*, 38, 3405-3415.

Hussein, T., Glytsos, T., Ondráček, J., Dohányosová, P., Ždímal, V., Hämeri, Lazaridis, M., Smolík, J., Kulmala, M., 2006. Particle size Characterization and emission rates during indoor activities in a house, *Atmospheric Environment*, 40, 4285-4307.

See, S.W., Balasubramaniana, R., 2006. Risk assessment of exposure to indoor aerosols associated with Chinese cooking, *Environmental Research*, 102, 197-204.